

Gantmakher–Krein theorem for 2-totally nonnegative operators in ideal spaces

Olga Y. Kushel, Petr P. Zabreiko
Department of mechanics and mathematics,
Belorussian State University,
Nezavisimosti sq., 4, 220050, Minsk, Belarus,
e-mail: kushel@mail.ru; zabreiko@bsu.by

August 2008

Abstract.

The tensor and exterior squares of a completely continuous non-negative linear operator A acting in the ideal space $X(\Omega)$ are studied. The theorem representing the point spectrum (except, probably, zero) of the tensor square $A \otimes A$ in the terms of the spectrum of the initial operator A is proved. The existence of the second (according to the module) positive eigenvalue λ_2 , or a pair of complex adjoint eigenvalues of a completely continuous non-negative operator A is proved under the additional condition, that its exterior square $A \wedge A$ is also nonnegative.

Keywords: Total positivity, Ideal spaces, Tensor products, Exterior products, Point spectrum.

2000 Mathematics Subject Classification: Primary 47B65 Secondary 47A80, 47B38, 46E30.

1 Introduction

This paper presents the results of our study of the spectrum of 2-totally-nonnegative operators in ideal spaces. The theory of total positivity is mainly based on the results of F.R. Gantmakher and M.G. Krein (see [1]), concerning the properties of the spectrum of k -totally primitive matrices (i.e. matrices, which are nonnegative and primitive along with their j -th compound matrices ($1 < j \leq k$) up to the order k). In the most important case $k = n$ k -totally primitive matrices are called *oscillatory*. In monograph [1] the following statement was proved: *if the matrix $\mathbf{A}$ of a linear operator A in the space $\mathbb{R}^n$ is k -totally primitive, then the operator A has k positive simple eigenvalues $0 < \lambda_k < \dots < \lambda_2 < \lambda_1$, with a positive eigenvector e_1 corresponding to the maximal*

eigenvalue λ_1 , and an eigenvector e_j , which has exactly $j - 1$ changes of sign, corresponding to j -th eigenvalue λ_j (see [1], p. 310, theorem 9).

The study of linear integral operators with analogous properties predates the study of oscillatory matrices. The main results concerning this problem were received by O.D. Kellog (see [2]). He proved the theorem about spectral properties of continuous symmetric totally nonnegative kernels. Later this theorem was generalized by F.R. Gantmakher for the non-symmetric case. This result one can find in monograph [1] in the following form: *let $k(t, s) \in C[0, 1]^2$ satisfy the following conditions:*

(a) *for any $0 < t_0 < t_1 < \dots < t_n < 1$ and $0 < s_0 < s_1 < \dots < s_n < 1$ $n = 0, 1, \dots$ the inequality*

$$k \begin{pmatrix} t_0 & t_1 & \dots & t_n \\ s_0 & s_1 & \dots & s_n \end{pmatrix} \geq 0$$

is true;

(b) *for any $0 < t_0 < t_1 < \dots < t_n < 1$ $n = 0, 1, \dots$ the inequality*

$$k \begin{pmatrix} t_0 & t_1 & \dots & t_n \\ t_0 & t_1 & \dots & t_n \end{pmatrix} > 0$$

is true.

Then all the eigenvalues of the linear integral equation

$$\int_0^1 k(t, s)x(s)ds = \lambda x(t)$$

are positive and simple:

$$0 < \dots < \lambda_n < \dots < \lambda_2 < \lambda_1,$$

with a strictly positive on $(0, 1)$ eigenfunction $e_1(t)$ corresponding to the maximal eigenvalue λ_1 , and an eigenfunction $e_n(t)$, which has exactly $n - 1$ changes of sign and no other zeroes on $(0, 1)$, corresponding to the n -th eigenvalue λ_n (see [1], p. 211).

The proof of this statement one can find also in [3], where the history of the theory of totally positive matrices and kernels is presented in detail. Unlike monograph [1], in which the basic statements of the theory are given in the form most suitable for the study of small oscillations of mechanical systems, in [3] definitions and theorems about the properties of totally positive kernels are given in the pure form.

In paper [4] by S.P. Eveson the result mentioned was spread onto a wider class of kernels. The existence of k positive eigenvalues was proved under some additional assumptions for the case of a compact linear integral operator, acting in $L_2[0, 1]$, which kernel is totally positive of order k . A substantial contribution into the development of the theory of totally positive and sign-symmetric kernels was made by S. Karlin (see [5]).

Once a great number of papers are devoted to the theory of totally positive matrices and kernels, in the case of abstract (not necessarily integral) compact linear operators the situation is absolutely different. Here we can mention only a small number of papers. In paper [6] oscillatory operators in $C[a, b]$ were studied by the method of the passage to the limit from finite-dimensional approximants. In paper [7] another method of generalization was suggested. But this method was realized also only for the space $C[a, b]$. Many results, related to the applications of the theory of oscillation to differential operators, were included into monograph [8] by Yu.V. Pokornyi and his group.

In paper [9] we studied 2-totally indecomposable operators (i.e. indecomposable operators that are nonnegative with respect to some cone K , and such that their exterior squares are also nonnegative and indecomposable) in the spaces $L_p(\Omega)$ ($1 \leq p \leq \infty$) and $C(\Omega)$. We proved the

existence of the second (according to the module) eigenvalue λ_2 of a completely continuous non-negative operator A under the condition that its exterior square $A \wedge A$ is also non-negative. The simplicity of the first and second eigenvalues was proved and the interrelation between the indices of imprimitivity of A and $A \wedge A$ was examined for the case, when the operators A and $A \wedge A$ are indecomposable. The difference (according to the module) of λ_1 and λ_2 from each other and from other eigenvalues was proved for the case, when A and $A \wedge A$ are primitive.

In the present paper we are going to generalize the results, received in [9], for 2-totally nonnegative operators in some ideal spaces. As the authors believe, the natural method of the examination of such operators is a crossway from studying an operator A , acting in an ideal space X to the study of the operators $A \otimes A$ and $A \wedge A$, acting in spaces with mixed norms. Let us turn now to a more detailed outline of the paper. In Section 2 we briefly consider the basic properties of ideal spaces. Tensor and exterior squares of ideal spaces are described in Section 3. The connection between the topological exterior square of the ideal space X and the space with mixed norm is also clarified in Section 3. In Section 4 we work with the tensor and exterior square of a linear operator $A : X \rightarrow X$. These operators act in the tensor and exterior square of the initial space X respectively. Generally, in Sections 3 and 4 we develop the mathematical tools that will enable us to define the class of abstract totally nonnegative operators and to generalize the results of Gantmakher and Krein. Sections 5 and 6 present a number of results on the description of the spectrum and the parts of the spectrum of the tensor square and the exterior square of a completely continuous non-negative linear operator A acting in the ideal space $X(\Omega)$ in terms of the spectrum of the initial operator. The main mathematical results of this paper are concentrated in Section 6, where we prove the existence of the second according to the module positive eigenvalue λ_2 , or a pair of complex adjoint eigenvalues of a completely continuous non-negative operator A under the additional condition, that its exterior square $A \wedge A$ is also nonnegative. For the case when A is a linear integral operator, the main theorem is formulated in terms of kernels.

2 Ideal spaces. Basic definitions and statements

Let $(\Omega, \mathfrak{A}, \mu)$ be a triple, consisting of a set Ω , σ -algebra $\mathfrak{A}$ of measurable subsets and σ -finite and σ -additive complete measure on $\mathfrak{A}$. Denote by $S(\Omega)$ the space of all measurable finite almost everywhere on Ω functions (further we consider the equivalent functions to be identical). Let $X(\Omega)$ be a Banach ideal space, i.e. a Banach space of all measurable on Ω functions having the following property: from $|x_1| \leq |x_2|$, $x_1 \in S(\Omega)$, $x_2 \in X$, it follows that $x_1 \in X$ $\|x_1\|_X \leq \|x_2\|_X$ (the definition and basic properties of ideal spaces are taken from paper [10], see also [11]). Consider the support $\text{supp} X$ of the space X to be the least measurable subset, outside which all the functions from X are equal to zero. Let $\text{supp} X = \Omega$, i.e. there exist functions from the space X , which are positive almost everywhere on Ω . The Banach ideal space X is called *regular*, if the norm in X is order-continuous, i.e. for every sequence $\{x_n\} \subset X$ from $0 \leq x_n \downarrow 0$ it follows that $\|x_n\| \rightarrow 0$. X is called *almost perfect*, if the norm in X is order-semicontinuous, i.e. for every sequence $\{x_n\} \subset X$ from $0 \leq x_n \uparrow x \in X$ it follows that $\|x_n\| \rightarrow \|x\|$. It's easy to see (see, for example, [10]), that every regular space is almost perfect.

Let us denote by $X_{12}(\Omega \times \Omega)$ the set of all measurable with respect to all the variables functions $x(t_1, t_2)$ on $\Omega \times \Omega$, which satisfy the following conditions:

- (1) for almost every t_2 the function $x(\cdot, t_2)$ belongs to X ;
- (2) the function $\|x(\cdot, t_2)\|$ also belongs to X .

If the space X is almost perfect then the set X_{12} is linear (see, for example [11], [12]). The norm in X_{12} is introduced according to the following rule:

$$\|x(t_1, t_2)\|_{12} = \| \|x(t_1, t_2)\|_{(1)} \|_{(2)},$$

where indices (1) and (2) mean, that the norm of the space X is used firstly for the first variable, and then for the second variable.

The space $X_{21}(\Omega \times \Omega)$ with the norm

$$\|x(t_1, t_2)\|_{21} = \| \|x(t_1, t_2)\|_{(2)} \|_{(1)}$$

is defined similarly.

In the case of an almost perfect X both the space X_{12} and the space X_{21} are almost perfect Banach ideal spaces (see [10], [11], and also [13], p. 1234, theorem 3, where the completeness of the space with mixed norm is proved).

Further we will be interested in the space $\tilde{X} = X_{12} \cap X_{21}$ of functions that are common for the spaces X_{12} and X_{21} . The norm in this space is introduced by the formula:

$$\|x(t_1, t_2)\|_M = \max\{\|x(t_1, t_2)\|_{12}, \|x(t_1, t_2)\|_{21}\}.$$

Note that the space $\tilde{X}$ is regular if and only if the space X is regular.

In connection with the introduced intersection of the spaces X_{12} and X_{21} there arises a natural question of the possibility of their coincidence. For $X = L_p$ from the Fubini theorem it follows that X_{12} and X_{21} coincide according to the fund of elements and according to their norms. However, this is not true in the general case. Moreover, the coincidence of the spaces X_{12} and X_{21} is characteristic for the class of the spaces L_p . For the regular Banach ideal space X N.J. Nielsen proved, that from $X_{12} = X_{21}$ it follows that X is lattice-isomorphic to L_p -space (see [14]). The results concerning this problem one can also find in [15] and [16].

3 Tensor and exterior squares of ideal spaces

The algebraic tensor square $X \otimes X$ of the space X is defined as the set of all functions of the form

$$x(t_1, t_2) = \sum_{i=1}^n x_1^i(t_1) x_2^i(t_2),$$

where $x_1^i, x_2^i \in X$. Further call the elements of $X \otimes X$ *degenerate functions*. By the way, the algebraic exterior square $X \wedge X$ of the space X is defined as the set of all antisymmetric functions (i.e. functions $x(t_1, t_2)$, for which $x(t_1, t_2) = -x(t_2, t_1)$) from $X \otimes X$.

Generally, the norm on $X \otimes X$ can be defined by different ways. Let us go through definitions, that will be used further. The norm α on $X \otimes X$ is called a *crossnorm*, if for any $x_1, x_2 \in X$ the following equality holds:

$$\|x_1(t_1)x_2(t_2)\|_\alpha = \|x_1(t_1)\| \|x_2(t_2)\|.$$

There exists the greatest crossnorm π (see, for example [17]), which is defined by the equality:

$$\|x(t_1, t_2)\|_\pi = \inf \sum_{i=1}^n \|x_1^i\| \|x_2^i\|,$$

where $\inf$ is taken over all representations $x(t_1, t_2) = \sum_{i=1}^n x_1^i(t_1)x_2^i(t_2)$, $x_1^i, x_2^i \in X$.

The least crossnorm does not exist, however there exists the least *reasonable* crossnorm (the norm α on $X \otimes X$ is called reasonable, if α is a crossnorm, and the dual norm α' is also a crossnorm on $X' \otimes X'$, where the dual norm is defined on $X' \otimes X'$ by the equality $\|x'\|_{\alpha'} = \sup \{\langle x, x' \rangle, \|x\|_\alpha \leq 1\}$, $x \in X \otimes X$, $x' \in X' \otimes X'$). The least reasonable crossnorm is denoted by ϵ and is defined by the following rule:

$$\|x\|_\epsilon = \sup \left\{ \left| \sum_{i=1}^n \langle x_1^i, x_1' \rangle \langle x_2^i, x_2' \rangle \right|, \|x_1'\|_{X'}, \|x_2'\|_{X'} \leq 1 \right\}.$$

Note that the completion of the algebraic tensor square $X \otimes X$ of the ideal space X with respect to the norms π or ϵ will not be ideal. It is natural to define the norm on $X \otimes X$ in such a way that the completion with respect to this norm will be an ideal space or its part. With this purpose V.L. Levin introduced in [18] the following crossnorm on $X \otimes X$:

$$\|x\|_L = \inf \left\{ \|u\|_X : u \geq \left| \sum_i x_1^i \langle x_2^i, x^* \rangle \right|, \|x^*\|_{X^*} \leq 1 \right\}.$$

V.L. Levin proved (see [18], p. 55, proof of the theorem 1), that the topology of the space X_{21} induces on $X \otimes X$ the same topology, as the norm L .

Note that every norm α on $X \otimes X$ which satisfies inequalities $\epsilon \leq \alpha \leq \pi$ will be reasonable. In particular, the Levin's norm L is reasonable (see [18], p. 53, lemma 3).

Further let us call the completion of $X \otimes X$ with respect to the Levin's norm the L -complete tensor square of the space X and let us denote it by $\widetilde{(X \otimes X)}_L$. As it was noticed above, $\widetilde{(X \otimes X)}_L$ is a closed subspace of X_{21} . The space $\widetilde{(X \otimes X)}_L$ was studied thoroughly by A.V. Bukhvalov in papers [19] and [13]. He also proved the criteria of the coincidence of this space and X_{21} (see [13], p. 7, theorem 0.1, and also [19], p. 1235, theorem 4).

Bukhvalov's theorem. *Let X be a Banach ideal space. Then the following statements are equivalent:*

- (i) *the set of all degenerate functions is dense in X_{21} ;*
- (ii) *the equality $\widetilde{(X \otimes X)}_L = X_{21}$ is true;*
- (iii) *the space X is regular.*

Bukhvalov's theorem implies that for a regular Banach ideal space the following equality holds:

$$\widetilde{(X \otimes X)}_L^a = X_{21}^a(\Omega \times \Omega),$$

where $X_{21}^a(\Omega \times \Omega)$ is a subspace of the space $X_{21}(\Omega \times \Omega)$, which consists of antisymmetric functions.

Further note that $\wedge$ -product of arbitrary functions $x_1, x_2 \in X$ ($x_1 \wedge x_2$)(t_1, t_2) = $x_1(t_1)x_2(t_2) - x_1(t_2)x_2(t_1)$ belongs to the space $X_{21}(\Omega \times \Omega)$, and the following equality holds:

$$x_1 \wedge x_2(t_1, t_2) = -x_1 \wedge x_2(t_2, t_1).$$

Therefore, for any $x_1, x_2 \in X$ the function $x_1 \wedge x_2$ belongs to the subspace $X_{21}^a(\Omega \times \Omega)$.

Moreover, an arbitrary antisymmetric function $x(t_1, t_2)$, which belongs to the space X_{21} , at the same time belongs to X_{12} , and the following equality holds

$$\|x(t_1, t_2)\|_{21} = \|x(t_1, t_2)\|_{12}.$$

Really, let $x(t_1, t_2)$ be equal to $-x(t_2, t_1)$. In this case

$$\begin{aligned} \|x(t_1, t_2)\|_{21} &= \| \|x(t_1, t_2)\|_{(2)} \|_{(1)} = \| \| -x(t_2, t_1) \|_{(2)} \|_{(1)} = \\ &= \| \|x(t_1, t_2)\|_{(1)} \|_{(2)} = \|x(t_1, t_2)\|_{12}. \end{aligned}$$

That is why further we will assume that the completion of the algebraic exterior square $X \wedge X$ of the space X is taken with respect to the symmetric crossnorm

$$\|x(t_1, t_2)\|_M = \max\{\|x(t_1, t_2)\|_{12}, \|x(t_1, t_2)\|_{21}\}.$$

This completion coincides with the closed subspace of antisymmetric functions of the space $\widetilde{X}(\Omega \times \Omega)$.

The subspace $\widetilde{X}^a(\Omega \times \Omega)$ is isomorphic in the category of Banach spaces to the space $\widetilde{X}(W)$, where W is the measurable subset $\Omega \times \Omega$, for which the sets $W \cap \widetilde{W}$ and $(\Omega \times \Omega) \setminus (W \cup \widetilde{W})$ have zero measure; here $\widetilde{W} = \{(t_2, t_1) : (t_1, t_2) \in W\}$ (such sets do always exist). Really, extending the functions from $\widetilde{X}(W)$ as antisymmetric functions to $\Omega \times \Omega$, we obtain the set of all the functions from $\widetilde{X}^a(\Omega \times \Omega)$. Further, setting the norm of a function in $\widetilde{X}(W)$ to be equal to the norm of its extension, we get that the spaces $\widetilde{X}^a(\Omega \times \Omega)$ and $\widetilde{X}(W)$ are isomorphic in the category of normed spaces.

Therefore, for an almost perfect Banach ideal space the following equality holds:

$$(\widetilde{X \wedge X})_M = (\widetilde{X \otimes X})_M^a = \widetilde{X}_d(W),$$

where $\widetilde{X}_d(W)$ is the closure of the set of all degenerate functions from $\widetilde{X}(W)$ in the norm of $\widetilde{X}(W)$.

4 Tensor and exterior squares of linear operators in ideal spaces

Let A, B be continuous linear operators acting in the ideal space X . Define the *algebraic tensor product* of the operators A and B as the operator $A \otimes B$ in the space $X \otimes X$, defined on degenerate functions by the equality

$$(A \otimes B)x(t_1, t_2) = \sum_j Ax_1^j(t_1) \cdot Bx_2^j(t_2) \quad \left(x(t_1, t_2) = \sum_j x_1^j(t_1) \cdot x_2^j(t_2) \right).$$

A crossnorm α on $X \otimes X$ is called *quasiuniform*, if for any continuous linear operators A, B one has:

$$\|A \otimes B\|_\alpha \leq c \|A\| \|B\|,$$

where c is some constant. If $c = 1$, then such crossnorm is called *uniform*.

The greatest and the least reasonable norms π and ϵ are uniform. However Levin's norm L in general case will not be quasiuniform. That is why we need the following definition.

Define the *tensor product* of the operators A and B (if it exists) as the linear operator $A \otimes B$ in the space $(\widetilde{X \otimes X})_L$, defined on degenerate functions by the equality

$$(A \otimes B)x(t_1, t_2) = \sum_j Ax_1^j(t_1) \cdot Bx_2^j(t_2) \quad \left(x(t_1, t_2) = \sum_j x_1^j(t_1) \cdot x_2^j(t_2) \right),$$

and on arbitrary functions by extension via continuity from the subspace of degenerate functions onto the whole of $(\widetilde{X \otimes X})_L$ (if the extension is bounded).

Let us formulate the following statement concerning with the tensor product of two operators in ideal spaces (see [18], p. 62, proposition 6).

Levin's theorem. *Let X be a Banach ideal space, $A : X \rightarrow X$ be a regular operator, i.e. an operator, which can be represented in the form $A = A_1 - A_2$, where A_1 and A_2 are nonnegative linear operators (with respect to the cone of nonnegative functions), and let $B : X \rightarrow X$ be a continuous linear operator. Then the tensor product $A \otimes B$ does exist.*

Further, when studying the spectral properties of the tensor square $A \otimes A$ of the operator A , we will have to impose conditions on the operator A , that are stronger than nonnegativity or regularity. Let us give the following definition. A linear operator $A : X \rightarrow X$ is called *resolvent-regular*, if for any λ , which is not in $\sigma(A)$, the resolvent operator $R(\lambda, A) = (\lambda I - A)^{-1}$ is regular.

The class of resolvent-regular operators includes, for example, Hilbert-Schmidt operators and operators, such that their certain power is a Hilbert-Schmidt operator.

Further let us examine the operator $A \wedge A$, defined as the restriction of the operator $A \otimes A$ onto the subspace $\widetilde{(X \otimes X)}_M^a$. It is obvious that for degenerate antisymmetric functions the operator $A \wedge A$ can be defined by the equality

$$(A \wedge A)x(t_1, t_2) = \sum_j Ax_1^j(t_1) \wedge Ax_2^j(t_2) \quad x(t_1, t_2) = \sum_j x_1^j(t_1) \wedge x_2^j(t_2).$$

5 Spectrum of the tensor square of linear operators in ideal spaces

As usual, we denote by $\sigma(A)$ the spectrum of an operator A , and we denote by $\sigma_p(A)$ the point spectrum, that is, the set of all eigenvalues of the operator A . Denote by $\sigma_{eb}(A)$ the Browder essential spectrum of the operator A . Thus $\sigma(A) \setminus \sigma_{eb}(A)$ will be the set of all isolated finite dimensional eigenvalues of the operator A (for more detailed information see [20], [21]).

In papers [20]–[23] T. Ichinose obtained the results, representing the spectra and the parts of the spectra of the tensor products of linear bounded operators in terms of the spectra and parts of the spectra of the operators given under the natural assumptions, that the corresponding crossnorm is reasonable and quasiuniform. Among the mentioned results there are the explicit formulae, expressing the set of all isolated finite dimensional eigenvalues and the Browder essential spectrum of the operator $A \otimes A$ in terms of the parts of the spectrum of the operator given (see [20], p. 110, theorem 4.2). In particular, Ichinose proved, that for the tensor square of a linear bounded operator A the following equalities hold:

$$\begin{aligned} \sigma(A \otimes A) &= \sigma(A) \sigma(A); \\ \sigma(A \otimes A) \setminus \sigma_{eb}(A \otimes A) &= (\sigma(A) \setminus \sigma_{eb}(A)) (\sigma(A) \setminus \sigma_{eb}(A)) \setminus (\sigma_{eb}(A) \sigma(A)); \\ \sigma_{eb}(A \otimes A) &= \sigma_{eb}(A) \sigma(A). \end{aligned} \tag{1}$$

For a completely continuous operator the following equalities hold:

$$(\sigma(A) \setminus \sigma_{eb}(A)) \setminus \{0\} = \sigma_p(A) \setminus \{0\}; \quad \sigma_{eb}(A) = \{0\} \text{ or } \emptyset.$$

So from (1) we can get the complete information about the nonzero eigenvalues of the tensor square of a completely continuous operator:

$$\sigma_p(A \otimes A) \setminus \{0\} = (\sigma_p(A) \sigma_p(A)) \setminus \{0\}. \tag{2}$$

Here zero can be either a finite or infinite dimensional eigenvalue of $A \otimes A$ or a point of the essential spectrum.

In paper [9] there have been examined the case, when a linear operator acts in the space $L_p(\Omega)$ ($C(\Omega)$), and the corresponding crossnorm is reasonable and quasiuniform. That is why the formula for the spectrum of $A \otimes A$ directly follows from the results of T. Ichinose.

However in general case the crossnorm L is reasonable, but not quasiuniform (see [24]), and therefore we need a different proof for the statement about the spectrum of $A \otimes A$. The proof, given below, is based on the reasoning, made by A.S. Kalitvin (see [11], p. 83, theorem 3.10), for the case of the operator $A \otimes I + I \otimes A$ in a regular ideal space.

Theorem 1. *Let X be an almost perfect Banach ideal space, and let $A : X \rightarrow X$ be a completely continuous nonnegative with respect to the cone of nonnegative functions in X resolvent-regular operator. Then for the point spectrum of the operator $A \otimes A$, acting in the space $\widetilde{(X \otimes X)}_L$, the following equality holds:*

$$\sigma_p(A \otimes A) \setminus \{0\} = (\sigma_p(A) \sigma_p(A)) \setminus \{0\}.$$

Proof. Let us examine the operators $A \otimes I$ and $I \otimes A$, acting in $\widetilde{(X \otimes X)}_L$. Let us prove, that the following inclusions are true: $\sigma(A \otimes I) \subseteq \sigma(A)$ and $\sigma(I \otimes A) \subseteq \sigma(A)$. Let us prove the first inclusion (the second inclusion can be proved by analogy). Let λ does not belong to $\sigma(A)$. Then the operator $\lambda I - A$ is invertible. Let us define the operator $(\lambda I - A)^{-1} \otimes I$ on $X \otimes X$. Since the operator $(\lambda I - A)^{-1}$ is regular, we can apply the Levin's theorem. As it follows from Levin's theorem, the operator $(\lambda I - A)^{-1} \otimes I$ can be extended from $X \otimes X$ onto the whole of $\widetilde{(X \otimes X)}_L$. It is easy to see, that the operator $(\lambda I - A)^{-1} \otimes I$ is inverse on $X \otimes X$ for the operator $\lambda I - (A \otimes I)$. So, its extension $(\lambda I - A)^{-1} \otimes I$ will be inverse for the operator $\lambda I - (A \otimes I)$ on the whole of $\widetilde{(X \otimes X)}_L$. That is why λ does not belong to $\sigma(A \otimes I)$, and the inclusion $\sigma(A \otimes I) \subseteq \sigma(A)$ is proved.

Thus as $A \otimes A = (A \otimes I)(I \otimes A)$ and the operators $A \otimes I$ and $I \otimes A$ are, obviously, commuting, the following relation is true

$$\sigma(A \otimes A) = \sigma((A \otimes I)(I \otimes A)) \subseteq \sigma(A \otimes I)\sigma(I \otimes A).$$

Now, applying the inclusions $\sigma(A \otimes I) \subseteq \sigma(A)$ and $\sigma(I \otimes A) \subseteq \sigma(A)$, proved above, we see that the following inclusion is true as well

$$\sigma(A \otimes A) \subseteq \sigma(A)\sigma(A).$$

since due to the complete continuity of the operator A , its spectrum, except, probably, zero, consists of isolated finite dimensional eigenvalues. That is why the following relations hold:

$$\begin{aligned} \sigma_p(A \otimes A) \setminus \{0\} &\subseteq \sigma(A \otimes A) \setminus \{0\} \subseteq (\sigma(A)\sigma(A)) \setminus \{0\} = \\ &= (\sigma_p(A)\sigma_p(A)) \setminus \{0\}, \end{aligned}$$

i.e. we proved:

$$\sigma_p(A \otimes A) \setminus \{0\} = (\sigma_p(A)\sigma_p(A)) \setminus \{0\}.$$

Now, let us prove the reverse inclusion. For this we will examine the extension $\widetilde{(A \otimes A)}_\epsilon$ of the operator $A \otimes A$ onto the whole $\widetilde{(X \otimes X)}_\epsilon$, where $\widetilde{(X \otimes X)}_\epsilon$ is a completion $X \otimes X$ with respect to the "weak" crossnorm ϵ . As it follows from the results of J.R. Holub (see [25], p. 401, theorem 2), the operator $\widetilde{(A \otimes A)}_\epsilon$ is completely continuous in $\widetilde{(X \otimes X)}_\epsilon$. Let us prove, that $\sigma_p(\widetilde{(A \otimes A)}_\epsilon) \setminus \{0\} \subseteq \sigma_p(A \otimes A) \setminus \{0\}$. To check this it is enough to prove, that any eigenfunction of the operator $\widetilde{(A \otimes A)}_\epsilon$, corresponding to a nonzero eigenvalue, belongs to the space $\widetilde{(X \otimes X)}_L$. Let λ be an arbitrary nonzero eigenvalue of the operator $\widetilde{(A \otimes A)}_\epsilon$. Formula (3) implies that there exist indices i, j , for which $\lambda = \lambda_i \lambda_j$ (here $\{\lambda_k\}$ is the set of all nonzero eigenvalues of the operator A , enumerated without regard to multiplicity). Since λ is an isolated finite dimensional eigenvalue, i, j can take only finite number of different values. Let us enumerate all the pairs of such values. Let $\lambda = \lambda_{i_k} \lambda_{j_k}$ ($k = 1, \dots, p$). Decompose the space X into the direct sum of subspaces:

$$X = X_1 \oplus \dots \oplus X_p \oplus R,$$

where $X_k = \ker(A - \lambda_{i_k})^{m_k}$, m_k are the multiplicities of λ_{i_k} . Under this decomposition $\widetilde{(X \otimes X)}_\epsilon$ also decomposes into the direct sum of subspaces:

$$\begin{aligned} \widetilde{(X \otimes X)}_\epsilon &= (X_1 \otimes X_1) \oplus \dots \oplus (X_1 \otimes X_p) \oplus \dots \oplus (X_p \otimes X_1) \oplus \dots \oplus (X_p \otimes X_p) \oplus \\ &\oplus (X_1 \otimes R) \oplus \dots \oplus (X_p \otimes R) \oplus (R \otimes X_1) \oplus \dots \oplus (R \otimes X_p) \oplus \widetilde{(R \otimes R)}_\epsilon. \end{aligned}$$

Since $X_l \otimes X_m$, $X_l \otimes R$, $R \otimes X_l$, $(\widetilde{R \otimes R})_\epsilon$ are invariant subspaces for the operator $(\widetilde{A \otimes A})_\epsilon$, the following equality holds:

$$\begin{aligned} & \sigma(\widetilde{A \otimes A})_\epsilon = \\ & = \bigcup_{l,m} (\sigma(A \otimes A, X_l \otimes X_m) \cup \sigma(A \otimes A, R \otimes X_m) \cup \sigma(A \otimes A, X_l \otimes R)) \cup \sigma(A \otimes A, \widetilde{R \otimes R}_\epsilon), \end{aligned}$$

where the notation $\sigma(A \otimes A, X_l \otimes X_m)$ means the spectrum of the restriction of the operator $A \otimes A$ onto the corresponding subspace. Since $X_l \otimes X_m$, $R \otimes X_m$, $X_l \otimes R$, $(\widetilde{R \otimes R})_\epsilon$ are the spaces with uniform crossnorms, we can apply the results of T. Ichinose, and therefore the following equalities hold:

$$\sigma(A \otimes A, X_l \otimes X_m) = \sigma(A, X_l) \sigma(A, X_m); \quad \sigma(A \otimes A, X_l \otimes R) = \sigma(A, X_l) \sigma(A, R);$$

$$\sigma(A \otimes A, R \otimes X_m) = \sigma(A, R) \sigma(A, X_m); \quad \sigma(A \otimes A, (\widetilde{R \otimes R})_\epsilon) = \sigma(A, R) \sigma(A, R).$$

Since λ_{i_k} and λ_{j_k} do not belong $\sigma(A, R)$ for any values of indices k ($k = 1, \dots, p$), λ does not belong to $\bigcup_{l,m} (\sigma(A \otimes A, R \otimes X_m) \cup \sigma(A \otimes A, X_l \otimes R)) \cup \sigma(A \otimes A, (\widetilde{R \otimes R})_\epsilon)$. As it follows, $\lambda \in \bigcup_{l,m} (\sigma(A \otimes A, X_l \otimes X_m))$. Further it is obvious, that for any l, m ($1 \leq l, m \leq p$) $X_l \otimes X_m$ belongs to the algebraic tensor square $X \otimes X$ and therefore it belongs to the space $(\widetilde{X \otimes X})_L$. So, for an arbitrary $\lambda \in \sigma_p((\widetilde{A \otimes A})_\epsilon) \setminus \{0\}$ the inclusion $\lambda \in \sigma_p(A \otimes A) \setminus \{0\}$ is true. $\square$

Let us notice, that under conditions of Theorem 1 the inclusion:

$$\sigma_{eb}(A \otimes A) \subseteq \sigma_{eb}(A) \sigma_{eb}(A) = \{0\}$$

follows from its proof. Moreover, for an arbitrary $\lambda \in (\sigma_p(A \otimes A) \setminus \{0\})$ the following equality holds:

$$\ker(A \otimes A - \lambda I \otimes I) = \bigoplus_k \ker(A - \lambda_{i_k} I) \otimes \ker(A - \lambda_{j_k} I),$$

where the summation is taken over all the numbers k of the pairs $\lambda_{i_k}, \lambda_{j_k}$, for which $\lambda_{i_k}, \lambda_{j_k} \in (\sigma_p(A) \setminus \{0\})$ and $\lambda = \lambda_{i_k} \lambda_{j_k}$.

As it follows from the results of the paper [26], the statements of Theorem 1 will be also true for the case, when the tensor square $A \otimes A$ of the operator A acts in the space $(\widetilde{X \otimes X})_M$.

6 Spectrum of the exterior square of linear operators in ideal spaces

For the exterior square, which is the restriction of the tensor square, the following inclusions hold:

$$\sigma(A \wedge A) \subset \sigma(A \otimes A)$$

$$\sigma_p(A \wedge A) \subset \sigma_p(A \otimes A).$$

Applying Theorem 1, we can prove the following statement:

Theorem 2. *Let X be an almost perfect ideal space and let $A : X \rightarrow X$ be a completely continuous nonnegative with respect to the cone of nonnegative functions in X resolvent-regular operator. Let $\{\lambda_i\}$ be the set of all nonzero eigenvalues of the operator A , repeated according to multiplicity. Then all the possible products of the type $\{\lambda_i \lambda_j\}$, where $i < j$, form the set of all the possible (except, probably, zero) eigenvalues of the exterior square of the operator $A \wedge A$, repeated according to multiplicity.*

The proof copies the proof of the corresponding statement from [9] (see [9], p. 12, Theorem 1).

7 Generalization of the Gantmakher–Krein theorems in the case of 2-totally nonnegative operators in ideal spaces

Let us prove some generalizations of the Gantmakher–Krein theorems in the case of operators in ideal spaces, using the generalized Krein–Rutman theorem (see [27]) about completely continuous operators, leaving invariant an almost reproducing cone K in a Banach space (for such operators the spectral radius $\rho(A)$ belongs to $\sigma_p(A)$).

Theorem 3. *Let $X(\Omega)$ be an almost perfect ideal space, and, respectively, let $\tilde{X}_d(W)$ be the closure of the set of all degenerate functions in intersection of the spaces with mixed norms. Let $A : X(\Omega) \rightarrow X(\Omega)$ be a completely continuous nonnegative with respect to the cone of nonnegative functions in X resolvent-regular operator, such that $\rho(A) > 0$. Let the exterior square $A \wedge A : \tilde{X}_d(W) \rightarrow \tilde{X}_d(W)$ be nonnegative with respect to the cone of nonnegative functions in $\tilde{X}_d(W)$, and $\rho(A \wedge A) > 0$. Then the operator A has a positive eigenvalue $\lambda_1 = \rho(A)$. Moreover, if there is only one eigenvalue on the spectral circle $|\lambda| = \rho(A)$, then the operator A has the second positive eigenvalue $\lambda_2 < \lambda_1$. If there is more than one eigenvalue on the spectral circle $|\lambda| = \rho(A)$, then either there is at least one pair of complex adjoint among them, or λ_1 is a multiple eigenvalue.*

Proof. Enumerate eigenvalues of a completely continuous operator A , repeated according to multiplicity, in order of decrease of their modules:

$$|\lambda_1| \geq |\lambda_2| \geq |\lambda_3| \geq \dots$$

Applying the generalized Krein–Rutman theorem to A we get: $\lambda_1 = \rho(A) > 0$. Now applying the generalized Krein–Rutman theorem to the operator $A \wedge A$ we get: $\rho(A \wedge A) \in \sigma_p(A \wedge A)$. (Note, that from $\sigma_{eb}(\wedge^2 A) \subseteq \sigma_{eb}(\widetilde{(\otimes^2 A)}_{M_2}) \subseteq \sigma_{eb}(A)\sigma_{eb}(A) = \{0\}$, it follows, that the operator $A \wedge A$ also satisfy the conditions of the generalized Krein–Rutman theorem).

As it follows from the statement of theorem 2, the exterior square of the operator A has no other nonzero eigenvalues, except for all the possible products of the form $\lambda_i \lambda_j$, where $i < j$. So, we conclude that $\rho(A \wedge A) > 0$ can be represented in the form of the product $\lambda_i \lambda_j$ with some values of the indices i, j , $i < j$. Thus, if there is only one eigenvalue on the spectral circle $|\lambda| = \rho(A)$, from the fact that eigenvalues are numbered in decreasing order it follows that $\rho(A \wedge A) = \lambda_1 \lambda_2$. Therefore $\lambda_2 = \frac{\rho(A \wedge A)}{\lambda_1} > 0$.

If there is m ($m \geq 2$) eigenvalues on the spectral circle $|\lambda| = \rho(A)$, then $\rho(A \wedge A) = \lambda_i \lambda_j$, where $1 \leq i < j \leq m$. So, both λ_i and λ_j are situated on the spectral circle $|\lambda| = \rho(A)$, and from the positivity of their product it follows, that λ_i, λ_j are either a pair of complex adjoint eigenvalues, or both are positive and coincide with $\rho(A)$. $\square$

It is well-known (see, for example, [25], p. 55, corollary from the proposition 2.1), that a linear integral operator A , acting in the Banach ideal space $X(\Omega)$, is nonnegative if and only if its kernel $k(t, s)$ is nonnegative almost everywhere on Ω . It is also well-known (see [9]), that the exterior power of a linear integral operator can be considered as a linear integral operator, acting in the space $\tilde{X}_d(W)$ with the kernel equal to the second associated to the kernel of the operator given. That is why it is not difficult to reformulate Theorem 3 in terms of kernels of linear integral operators. In this case the conditions of Theorem 3 can be easily verified.

Theorem 4. *Let a completely continuous resolvent-regular linear integral operator A act in an almost perfect ideal space $X(\Omega)$. Let the kernel $k(t, s)$ of the operator A be nonnegative and not equal identically to zero almost everywhere on the Cartesian square $\Omega \times \Omega$. Let the second associated kernel $k \wedge k(t_1, t_2, s_1, s_2)$ be nonnegative and not equal identically to zero almost everywhere on the Cartesian square $W \times W$, where W is a measurable subset, possessing the following properties:*

- 1) $\mu(W \cap \widetilde{W}) = 0$;

$$2) \mu((\Omega \times \Omega) \setminus (W \cup \widetilde{W})) = 0. \quad (\widetilde{W} = \{(t_2, t_1) : (t_1, t_2) \in W\})$$

Then the operator A has a positive eigenvalue $\lambda_1 = \rho(A)$. Moreover, if there is only one eigenvalue on the spectral circle $|\lambda| = \rho(A)$, then the operator A has the second positive eigenvalue $\lambda_2 < \lambda_1$. If there is more than one eigenvalue on the spectral circle $|\lambda| = \rho(A)$, then either there is at least one pair of complex adjoint among them, or λ_1 is a multiple eigenvalue.

Note that in Theorem 4 the kernel is not presupposed to be continuous, we assume only, that the operator A acts in one of almost perfect ideal spaces.

Moreover, Theorem 3 can be generalized in the case, when the exterior square $A \wedge A$ of the operator A leaves invariant an arbitrary almost reproducing cone $\widetilde{K}$ in $\widetilde{X}_d(W)$. But in this case certain difficulties, related to the testing of the assumption of the generalized theorem, can arise.

References

1. *Gantmacher F.R., Krein M.G.* Oscillation Matrices and Kernels and Small Vibrations of Mechanical Systems. — AMS Bookstore, 2002. — 310 p.
2. *Kellog O.D.* Orthogonal function sets arising from integral equations // American Journal of Mathematics. — 1918. — Vol. 40. — P. 145-154.
3. *Pinkus A.* Spectral properties of totally positive kernels and matrices // Total positivity and its applications / M. Gasca, C.A. Micchelli. — Dordrecht, Boston, London: Kluwer Acad. Publ., 1996. — P. 1-35.
4. *Karlin S.* Total positivity. — Stanford University Press, California, 1968. — Vol. 1. — 576 p.
5. *Eveson S.P.* Eigenvalues of totally positive integral operators // Bull. London Math. Soc. — 1997. — Vol. 29. — P. 216-222.
6. *Sobolev A.V.* Abstract oscillatory operators // Proceedings of the Seminar on differential equations. — Kuibishev, 1977. — No. 3. — P. 72-78 (Russian).
7. *Yudovich V.I.* Spectral properties of an evolution operator of a parabolic equation with one space variable and its finite-dimensional analogues // Uspekhi Mat. Nauk. — 1977. — Vol. 32, No. 1. — P. 230-232 (Russian).
8. *Pokornyi Yu.V., Penkin O.M., Pryadiev V.L., Borovskikh A.V., Lazarev K.P., Shabrov S.A.* Differential equations on geometrical graphs. — Moscow: FIZMATLIT, 2004. — 272 p.
9. *Kushel O.Y., Zabreiko P.P.* Gantmakher-Krein theorem for 2-nonnegative operators in spaces of functions // Abstract and Applied Analysis. — 2006. — Article ID 48132. — P. 1-15.
10. *Zabreiko P.P.* Ideal spaces of functions // Vestnik Yaroslav. Univ. — 1974. — No. 4. — P. 12-52 (Russian).
11. *Kalitvin A.S.* Linear operators with partial integrals. — Voronezh, 2000. — 252 p. (Russian).
12. *Kantorovich L.V., Akilov G.P.* Functional Analysis. — 2nd rev. Moscow, 1977 (Russian). English transl.: Pergamon Press, Oxford, 1982.
13. *Bukhvalov A.V.* On spaces with mixed norm // Vestnik Leningrad. Univ. — 1973. — No. 19. — P. 512 (Russian); English transl.: Vestnik Leningrad Univ. Math. — 1979. — No. 6. — P. 303-311.
14. *Nielsen N.J.* On Banach ideals determined by Banach lattices and their applications // Diss. Math. — 1973. — 109.
15. *Bukhvalov A.V.* Generalization of the Kolmogorov-Nagumo theorem on the tensor product // Qualitative and Approximate Methods for the Investigation of Operator Equations. — Jaroslavl, 1979. — No. 4. — P. 4865 (Russian).
16. *Boccuto A., Bukhvalov A.V., Sambucini A.R.* Some inequalities in classical spaces with mixed norms // Positivity. — 2002. — Vol. 6. No. 4. — P. 393-411.
17. *Tsoy-Wo Ma.* Classical analysis on normed spaces. — World Scientific Publishing, 1995.
18. *Levin V.L.* Tensor products and functors in categories of Banach spaces defined by KB-lineals // Trudy Moskov. Mat. Obshch. — 1969. — Vol. 20. — P. 4381 (Russian); English transl.: Trans. Moscow Math. Soc. — 1969. — Vol. 20. — P. 4178.
19. *Bukhvalov A.V.* Vector-valued function spaces and tensor products // Siberian Math. J. — 1972. — Vol. 13, No. 6. — P. 1229-1238 (Russian).
20. *Ichinose T.* Spectral properties of tensor products of linear operators. I // Transactions of the American Mathematical Society. — 1978. — Vol. 235. — P. 75-113.
21. *Ichinose T.* Spectral properties of tensor products of linear operators. II // Transactions of the American Mathematical Society. — 1978. — Vol. 237. — P. 223-254.
22. *Ichinose T.* Operators on tensor products of Banach spaces // Trans. Am. Math. Soc. — 1972. — Vol. 170. — P. 197-219.

23. *Ichinose T.* Operational calculus for tensor products of linear operators in Banach spaces. // Hokkaido Math. J. – 1975. – Vol. 4. No. 2. – P. 306-334.
24. *Bukhvalov A.V.* Application of methods of the theory of order-bounded operators to the theory of operators in L^p -spaces // Uspekhi Mat. Nauk. – 1983. – Vol. 38, Issue 6(234). – P. 3783.
25. *Holub J.R.* Compactness in topological tensor products and operator spaces // Proc. Am. Math. Soc. – 1972. – Vol. 36. – P. 398-406.
26. *Zabreiko P.P.* The spectrum of linear operators acting in various Banach spaces // Qualitative and Approximate Methods for the Investigation of Operator Equations. – Jaroslavl, 1976. – No. 1. – P. 39-47 (Russian).
27. *Zabreiko P.P. and Smitskikh S.V.* A theorem of M. G. Krein and M. A. Rutman // Functional Analysis and Its Applications. – 1980. – Vol. 13. – P. 222-223.